
Index

- A-optimality, 439
- absolute error, 484, 494, 526
- ACM Transactions on Mathematical Software*, 552, 617
- $\text{adj}(\cdot)$, 69
- adjacency matrix, 331–334, 390
 - Exercise 8.20.*, 396
 - augmented, 390
- adjoint (see also conjugate transpose), 59
- adjoint, classical (see also adjugate), 69
- adjugate, 69, 598
- adjugate and inverse, 118
- affine group, 115, 179
- affine space, 43
- affine transformation, 228
- Aitken’s integral, 217
- $\mathcal{AL}(\cdot)$ (affine group), 115
- algebraic multiplicity, 144
- algorithm, 264, 499–513
 - batch, 512
 - definition, 509
 - direct, 264
 - divide and conquer, 506
 - greedy, 506
 - iterative, 264, 508–511, 564
 - reverse communication, 564
 - online, 512
 - out-of-core, 512
 - real-time, 512
- algorithmically singular, 121
- Anaconda, 553, 556, 569
- $\text{angle}(\cdot, \cdot)$, 37
- angle between matrices, 168
- angle between vectors, 37, 229, 357
- ANSI (standards), 475, 562, 563
- Applied Statistics* algorithms, 617
- approximation and estimation, 431
- approximation of a matrix, 174, 257, 339, 437, 533
- approximation of a vector, 41–43
- arithmetic mean, 35, 37
- Arnoldi method, 318
- artificial ill-conditioning, 269
- ASCII code, 460
- association matrix, 328–338, 357, 365, 366, 369–371
 - adjacency matrix, 331–332, 390
 - connectivity matrix, 331–332, 390
 - dissimilarity matrix, 369
 - distance matrix, 369
 - incidence matrix, 331–332, 390
 - similarity matrix, 369
- ATLAS (Automatically Tuned Linear Algebra Software), 555
- augmented adjacency matrix, 390
- augmented connectivity matrix, 390
- Automatically Tuned Linear Algebra Software (ATLAS), 555
- autoregressive process, 447–450
- axpy, 12, 50, 83, 554, 555
- axpy elementary operator matrix, 83
- back substitution, 274, 406
- backward error analysis, 495, 500
- Banach space, 33
- Banachiewicz factorization, 255

- banded matrix, 58
 - inverse, 121
- Bartlett decomposition, 346
- base, 467
- base point, 466
- basis, 21–23
 - Exercise 2.6*, 52
 - orthonormal, 40–41
- batch algorithm, 512
- Bauer-Fike theorem, 307
- Beowulf (cluster computing), 559
- bias, in exponent of floating-point number, 468
- big endian, 480
- big integer, 466, 492
- big O (order), 497, 503, 591
- big omega (order), 497
- bilinear form, 91, 134
- bit, 460
- bitmap, 461
- BLACS (software), 557, 558
- BLAS (software), 553–556
 - CUDA, 560
 - PBLAS, 558
 - PLASMA, 559
 - PSBLAS, 558
- block diagonal matrix, 62
 - determinant of, 70
 - inverse of, 121
 - multiplication, 78
- BMvN distribution, 219, 548
- Bolzano-Weierstrass theorem for orthogonal matrices, 133
- Boolean matrix, 390
- Box M statistic, 368
- bra-ket notation, 24
- byte, 460

- C (programming language), 474, 489, 568–569
- C++ (programming language), 475, 568–569
- CALGO (Collected Algorithms of the ACM)*, 617
- cancellation error, 487, 500
- canonical form, equivalent, 110
- canonical form, similar, 149
- canonical singular value factorization, 162

- Cartesian geometry, 35, 73
- catastrophic cancellation, 486
- Cauchy matrix, 389
- Cauchy-Schwarz inequality, 24, 98
- Cauchy-Schwarz inequality for matrices, 98, 178
- Cayley multiplication, 75, 94
- Cayley-Hamilton theorem, 138
- CDF (Common Data Format), 463
- centered matrix, 288, 363
- centered vector, 49
- chaining of operations, 485
- character data, 461
- character string, 461
- characteristic equation, 138
- characteristic polynomial, 138
- characteristic value (see also eigenvalue), 135
- characteristic vector (see also eigenvector), 135
- chasing, 317
- Chebyshev norm, 28
- chi-squared distribution, 400
 - noncentral, 400
 - PDF, equation (9.3), 400
- Cholesky decomposition, 253–256, 345, 436
 - computing, 558, 574
 - root-free, 255
- circulant matrix, 383
- classification, 389
- cluster analysis, 389
- cluster computing, 559
- coarray (Fortran construct), 563
- Cochran's theorem, 353–356, 401
- cofactor, 68, 598
- Collected Algorithms of the ACM (CALGO)*, 617
- collinearity, 265, 406, 430
- column rank, 99
- column space, 55, 89, 104, 105
- column-major, 522, 539, 545
- column-sum norm, 165
- Common Data Format (CDF), 463
- companion matrix, 139, 305
- compatible linear systems, 106
- compensated summation, 486
- complementary projection matrix, 356
- complementary vector spaces, 19

- complete graph, 329
- complete pivoting, 275
- complete space, 33
- completing the Gramian, 177
- complex data type, 491
- complex vectors/matrices, 33, 132, 387
- Conda, 553
- condition (problem or data), 499
- condition number, 265, 270, 290, 426, 427, 499, 502, 523, 533
 - computing the number, 533, 574
 - inverse of matrix, 266, 270
 - nonfull rank matrices, 290
 - nonsquare matrices, 290
 - sample standard deviation, 502
- conditional inverse, 128
- cone, 43–46, 327
 - Exercise 2.18.*; 53
 - convex cone, 44, 346
 - of nonnegative definite matrices, 346
 - of nonnegative matrices, 371
 - of positive definite matrices, 349
 - of positive matrices, 371
- conference matrix, 381
- configuration matrix, 369
- conjugate gradient method, 279–283
 - preconditioning, 282
- conjugate norm, 93
- conjugate transpose, 59, 132
- conjugate vectors, 93, 134
- connected vertices, 329, 334
- connectivity matrix, 331–334, 390
 - Exercise 8.20.*; 396
 - augmented, 390
- consistency property of matrix norms, 164
- consistency test, 527, 551
- consistent system of equations, 105, 272, 277
- constrained least squares, equality
 - constraints, 413
 - Exercise 9.4d.*; 451
- continuous function, 186
- contrast, 410
- convergence criterion, 508
- convergence of a sequence of matrices, 134, 152, 171
- convergence of a sequence of vectors, 32
- convergence of powers of a matrix, 171, 375
- convergence rate, 509
- convex combination, 12
- convex cone, 44–46, 346, 349, 371
- convex function, 26, 197
- convex optimization, 346
- convex set, 12
- convexity, 26
- coordinate, 4
- Cor(\cdot, \cdot), 51
- correlation, 51, 90
- correlation matrix, 365, 422
 - Exercise 8.8.*; 394
 - positive definite approximation, 436
 - pseudo-correlation matrix, 437
 - sample, 422
- cost matrix, 369
- Cov(\cdot, \cdot), 50
- covariance, 50
- covariance matrix, *see* variance-covariance matrix
- CRAN (Comprehensive R Archive Network), 552, 576
- cross product of vectors, 47
 - Exercise 2.19.*; 54
- cross products matrix, 256, 358
- cross products, computing sum of
 - Exercise 10.18c.*; 519
- Crout method, 245
- cuBLAS, 560
- CUDA, 559
- curl, 192
- curse of dimensionality, 511
- cuSPARSE, 560
- D-optimality, 439–440, 533
- daxpy**, 12
- decomposable matrix, 373
- decomposition, *see also* factorization of a matrix
 - additive, 353, 354
 - Bartlett decomposition, 346
 - multiplicative, 108
 - nonnegative matrix factorization, 257, 337
 - singular value decomposition, 161–164, 320, 336, 425, 532
 - spectral decomposition, 155

- defective (deficient) matrix, 149, 150
- deficient (defective) matrix, 149, 150
- deflation, 308–310
- degrees of freedom, 360, 362, 407, 430
- del, 192
- derivative with respect to a matrix, 195
- derivative with respect to a vector, 189–195
- $\det(\cdot)$, 66
- determinant, 66–74
 - as criterion for optimal design, 439
 - computing, 533
 - derivative of, 195
 - Jacobian, 217
 - of block diagonal matrix, 70
 - of Cayley product, 88
 - of diagonal matrix, 70
 - of elementary operator matrix, 86
 - of inverse, 117
 - of Kronecker product, 96
 - of nonnegative definite matrix, 345
 - of partitioned matrix, 70, 122
 - of permutation matrix, 87
 - of positive definite matrix, 347
 - of transpose, 70
 - of triangular matrix, 70
 - relation to eigenvalues, 141
 - relation to geometric volume, 73, 217
- $\text{diag}(\cdot)$, 56, 60, 596
 - with matrix arguments, 62
- diagonal element, 56
- diagonal expansion, 73
- diagonal factorization, 148, 152
- diagonal matrix, 57
 - determinant of, 70
 - inverse of, 120
 - multiplication, 78
- diagonalizable matrix, 148–152, 306, 343
 - orthogonally, 154
 - unitarily, 147, 343, 387
- diagonally dominant matrix, 57, 62, 100, 348
- differential, 188
- differentiation of vectors and matrices, 183–220
- digraph, 332
 - of a matrix, 333
- $\dim(\cdot)$, 14
- dimension of vector space, 14
- dimension reduction, 20, 356, 425
- direct method for solving linear systems, 272–277
- direct product (of matrices), 95
- direct product (of sets), 5
- direct product (of vector spaces), 20
 - basis for, 23
- direct sum decomposition of a vector space, 19
- direct sum of matrices, 62
- direct sum of vector spaces, 18–20, 64
 - basis for, 22
 - direct sum decomposition, 19
- directed dissimilarity matrix, 369
- direction cosines, 38, 231
- discrete Fourier transform, 384
- discrete Legendre polynomials, 380
- discretization error, 498, 509
- dissimilarity matrix, 369
- distance, 32
 - between matrices, 175
 - between vectors, 32
- distance matrix, 369
- distributed computing, 463, 508, 544
- distributed linear algebra machine, 558
- distribution vector, 377
- div, 192
- divergence, 192
- divide and conquer, 506
- document-term matrix, 336
- dominant eigenvalue, 142
- Doolittle method, 245
- dot product of matrices, 97
- dot product of vectors, 23, 91
- double cone, 43
- double precision, 472, 480
- doubly stochastic matrix, 377
- Drazin inverse, 129–130
- dual cone, 44
- E_{pq} , $E_{(\pi)}$, $E_p(a)$, $E_{pq}(a)$ (elementary operator matrices), 84
- $E(\cdot)$ (expectation operator), 216
- E-optimality, 439
- echelon form, 111
- edge of a graph, 329
- EDP (exact dot product), 493
- effective degrees of freedom, 362, 430

- efficiency, computational, 503–508
- eigenpair, 134
- eigenspace, 144
- eigenvalue, 134–164, 166, 305–322
 - computing, 306–318, 558, 574
 - Jacobi method, 313–316
 - Krylov methods, 318
 - power method, 311–313
 - QR method, 316–318
 - of a graph, 391
 - of a polynomial *Exercise 3.26*:, 181
 - relation to singular value, 163
 - upper bound on, 142, 145, 306
- eigenvector, 134–164, 305–322
 - left eigenvector, 135, 158
- eigenvectors, linear independence of, 143
- EISPACK, 556
- elementary operation, 79
- elementary operator matrix, 79–87, 101, 242, 273
 - eigenvalues, 137
- elliptic metric, 94
- elliptic norm, 93
- endian, 480
- equivalence of norms, 29, 33, 170
- equivalence relation, 444
- equivalent canonical factorization, 112
- equivalent canonical form, 110, 112
- equivalent matrices, 110
- error bound, 496
- error in computations
 - cancellation, 487, 500
 - error-free computations, 493
 - measures of, 484, 494–497, 526
 - rounding, 487, 495
 - Exercise 10.10*:, 517
- error of approximation, 498
 - discretization, 498
 - truncation, 498
- error, measures of, 272, 484, 494–497, 526
- error-free computations, 493
- errors-in-variables, 405
- essentially disjoint vector spaces, 15, 64
- estimable combinations of parameters, 409
- estimation and approximation, 431
- Euclidean distance, 32, 369
- Euclidean distance matrix, 369
- Euclidean matrix norm (see also Frobenius norm), 167
- Euclidean vector norm, 27
- Euler’s constant *Exercise 10.2*:, 516
- Euler’s integral, 593
- Euler’s rotation theorem, 231
- exact computations, 493
- exact dot product (EDP), 493
- exception, in computer operations, 483, 487
- expectation, 212–220
- exponent, 467
- exponential order, 503
- exponential, matrix, 153, 184
- extended precision, 472
- extrapolation, 509
- factorization of a matrix, 108, 112, 147, 148, 161, 225–227, 239–259, 272, 274
 - Banachiewicz factorization, 255
 - Bartlett decomposition, 346
 - canonical singular value factorization, 162
 - Cholesky factorization, 253–256
 - diagonal factorization, 148
 - equivalent canonical factorization, 112
 - full rank factorization, 108, 112
 - Gaussian elimination, 272
 - LQ factorization, 247
 - LU or LDU factorization, 240–246
 - nonnegative matrix factorization, 257, 337
 - orthogonally diagonal factorization, 147
 - QL factorization, 247
 - QR factorization, 246–252
 - root-free Cholesky, 255
 - RQ factorization, 247
 - Schur factorization, 147
 - singular value factorization, 161–164, 320, 336, 425, 532
 - square root factorization, 160
 - unitarily diagonal factorization, 147
- fan-in algorithm, 485, 506
- fast Fourier transform (FFT), 386
- fast Givens rotation, 239, 525

- fill-in, 259, 526
- Fisher information, 205
- fixed-point representation, 465
- flat, 43
- floating-point representation, 466
- FLOP, or flop, 505
- FLOPS, or flops, 505
- Fortran, 475–478, 505, 546, 566–568
- Fourier coefficient, 41, 42, 99, 157, 163, 169
- Fourier expansion, 36, 41, 98, 157, 163, 169
- Fourier matrix, 384
- Frobenius norm, 167–169, 171, 175, 314, 340, 369
 - Frobenius p norm, 169
- full precision, 480
- full rank, 100, 101, 104, 111–113
- full rank factorization, 108
 - symmetric matrix, 112
- full rank partitioning, 104, 122

- g_1 inverse, 128, 130
- g_2 inverse, 128
- g_4 inverse (see also Moore-Penrose inverse), 128
- gamma function, 220, 593
- GAMS (*Guide to Available Mathematical Software*), 539
- Gauss (software), 570
- Gauss-Markov theorem, 411
- Gauss-Newton method, 203
- Gauss-Seidel method, 277
- Gaussian elimination, 84, 272, 317
- Gaussian matrix, 84, 241
- gemm (general matrix-matrix), 556
- gemv (general matrix-vector), 556
- general linear group, 114, 133
- generalized eigenvalue, 160, 319
- generalized inverse, 124–125, 127–131, 249, 359
 - relation to QR factorization, 249
- generalized least squares, 414
- generalized least squares with equality constraints *Exercise 9.4d*, 451
- generalized variance, 366
- generating set, 14, 21
 - of a cone, 44
- generation of random numbers, 440

- geometric multiplicity, 144
- geometry, 35, 73, 227, 231
- Gershgorin disks, 145
- GitHub, 539
 - Exercise 12.3*., 581
- Givens transformation (rotation), 236–239, 317
 - QR factorization, 251
- $GL(\cdot)$ (general linear group), 114
- GMP (software library), 466, 491, 492
 - Exercise 10.5*., 516
- GMRES, 282
- GNU Scientific Library (GSL), 556
- GPU (graphical processing unit), 559
- graceful underflow, 470
- gradient, 189
 - projected gradient, 207
 - reduced gradient, 207
- gradient descent, 198, 199
- gradient of a function, 190, 191
- gradual underflow, 470, 487
- Gram-Schmidt transformation, 39, 40, 524
 - linear least squares, 289
 - QR factorization, 252
- Gramian matrix, 115, 117, 256, 289, 358–359
 - completing the Gramian, 177
- graph of a matrix, 332
- graph theory, 8, 329–336, 389
- graphical processing unit (GPU), 559
- greedy algorithm, 506
- group, 114, 133
- GSL (GNU Scientific Library), 556
- guard digit, 485

- Haar distribution, 219, 549
 - Exercise 4.10*., 221
 - Exercise 8.8*., 394
- Haar invariant measure, 220
- Hadamard matrix, 380
- Hadamard multiplication, 94
- Hadamard's inequality *Exercise 5.5*., 260, 606
- Hadoop, 464, 544
- Hadoop Distributed File System (HDFS), 464, 514
- half precision, 480
- Hankel matrix, 388

- Hankel norm, 389
- hat matrix, 360, 408
- HDF, HDF5 (Hierarchical Data Format), 463
- HDFS (Hadoop Distributed File System), 464, 514
- Helmert matrix, 378, 410
- Hemes formula, 286, 415
- Hermite form, 111
- Hermitian matrix, 56, 60
- Hessenberg matrix, 59, 316
- Hessian matrix, 193
 - projected Hessian, 207
 - reduced Hessian, 207
- Hessian of a function, 193
- hidden bit, 468
- Hierarchical Data Format (HDF), 463
- high-performance computing, 507
- Hilbert matrix, 548
- Hilbert space, 33, 168
- Hilbert-Schmidt norm (see also Frobenius norm), 167
- Hoffman-Wielandt theorem, 339
- Hölder norm, 27
- Hölder's inequality *Exercise 2.11a*, 52
- hollow matrix, 57, 369
- homogeneous coordinates, 232
 - in graphics applications, *Exercise 5.2*, 259
- homogeneous system of equations, 43, 123
- Horner's method, 512
- Householder transformation (reflection), 233–236, 250, 317
- hyperplane, 43
- hypothesis testing, 408

- idempotent matrix, 350–357
- identity matrix, 60, 76
- IDL (software), 6, 570
- IEC standards, 464
- IEEE standards, 464, 487
 - Standard 754, 472, 480, 487, 494
 - Standard P1788, 493
- IFIP Working Group 2.5, 460, 494
- ill-conditioned (problem or data), 264, 426, 499, 523
 - artificial, 269
 - stiff data, 502
- ill-posed problem, 121
- image data, 461
- IMSL Libraries, 556, 560–562
- incidence matrix, 331–334, 390
- incomplete data, 435–438
- incomplete factorization, 258, 526
- independence, linear, *see* linear independence
- independent vertices, 390
- index-index-value (sparse matrices), 548
- induced matrix norm, 165
- infinity, floating-point representation, 473, 487, 488
- infix operator, 489
- inner product, 23, 245
 - inner product of matrices, 96–99
 - inner product space, 24
- inner pseudoinverse, 128
- integer representation, 465
- integration and expectation, 212–220
- integration of vectors and matrices, 213
- Intel Math Kernel Library (MKL), 555, 578
- intersection graph, 334
- intersection of vector spaces, 18
- interval arithmetic, 492, 493
- invariance property, 227
- invariant distribution, 445
- invariant vector (eigenvector), 135
- inverse of a matrix, 107
 - determinant of, 117
 - Drazin inverse, 129–130
 - generalized inverse, 124–125, 127–131
 - Drazin inverse, 129–130
 - Moore-Penrose inverse, 127–129
 - pseudoinverse, 128
 - Kronecker product, 118
 - left inverse, 108
 - Moore-Penrose inverse, 127–129
 - partitioned matrix, 122
 - products or sums of matrices, 118
 - pseudoinverse, 128
 - right inverse, 108
 - transpose, 107
 - triangular matrix, 121
- inverse of a vector, 31
- IRLS (iteratively reweighted least squares), 297
- irreducible Markov chain, 444

- irreducible matrix, 311, 335–336, 372–377, 445
- `is.na`, 473
- `is.nan`, `is.nan`, 473
- ISO (standards), 475, 562, 563
- isometric matrix, 167
- isometric transformation, 227
- isotropic transformation, 228
- iterative method, 277, 284, 305, 508–511, 525
 - for solving linear systems, 277–283
 - iterative refinement, 284
- iteratively reweighted least squares, 297

- Jacobi method for eigenvalues, 313–316
- Jacobi transformation (rotation), 236
- Jacobian, 191, 217
- Jordan block, 77, 139
- Jordan decomposition, 151
- Jordan form, 77, 111
 - of nilpotent matrix, 77

- Kalman filter, 502
- Kantorovich inequality, 350
- Karush-Kuhn-Tucker conditions, 210
- kind (for data types), 476
- Kronecker multiplication, 94–96
 - inverse, 118
 - properties, 95
 - symmetric matrices, 96, 156
 - diagonalization, 156
- Kronecker structure, 219, 419
- Krylov method, 281, 318
- Krylov space, 281
- Kuhn-Tucker conditions, 210
- Kulisch accumulator, 493
- Kullback-Leibler divergence, 175

- L_1 , L_2 , and L_∞ norms
 - of a matrix, 165
 - of a symmetric matrix, 167
 - of a vector, 27
 - relations among, 170
- L_2 norm of a matrix (see also spectral norm), 166
- Lagrange multiplier, 208, 414
 - Exercise 9.4a.*, 450
- Lagrangian function, 208
- Lanczos method, 318

- LAPACK, 276, 534, 556, 558
 - LAPACK95, 556
- Laplace expansion, 68
- Laplace operator (∇^2), 599
- Laplace operator (∇^2), 192
- Laplacian matrix, 391
- lasso regression, 430
- latent root (see also eigenvalue), 135
- LAV (least absolute values), 295
- LDU factorization, 240–246
- leading principal submatrix, 62, 348
- least absolute values, 295
- least squares, 200–204, 256, 286–294
 - constrained, 206–211
 - nonlinear, 202–204
- least squares regression, 201
- left eigenvector, 135, 158
- left inverse, 108
- length of a vector, 4, 27, 31
- Leslie matrix, 378, 446
 - Exercise 8.10.*, 395
 - Exercise 9.22.*, 455
- Levenberg-Marquardt method, 204
- leverage, 408
 - Exercise 9.6.*, 451
- life table, 447
- likelihood function, 204
- line, 43
- linear convergence, 509
- linear estimator, 409
- linear independence, 12, 99
- linear independence of eigenvectors, 143
- linear programming, 346
- linear regression, 401–422, 426–431
 - variable selection, 427
- LINPACK, 276, 533, 556
- Lisp-Stat (software), 570
- little endian, 480
- little o (order), 497, 591
- little omega (order), 497
- log order, 503
- log-likelihood function, 205
- Longley data *Exercise 9.10.*, 453
- loop unrolling, 565
- Lorentz cone, 44
- lower triangular matrix, 57
- L_p norm
 - of a matrix, 165
 - of a vector, 27–28, 186

- LQ factorization, 247
- LR method, 306
- LU factorization, 240–246
 - computing, 558, 574
- M-matrix, 393
- MACHAR, 478
 - Exercise 10.3(d)i*, 516
- machine epsilon, 470
- Mahalanobis distance, 94, 365
- Manhattan norm, 27
- manifold of a matrix, 55
- Maple (software), 491, 570
- MapReduce, 463, 513, 531, 544
- Markov chain, 443–445
- Markov chain Monte Carlo (MCMC), 445
- Mathematica (software), 491, 570
- Matlab (software), 546, 578–580
- matrix, 5
- matrix derivative, 183–220
- matrix exponential, 153, 184
- matrix factorization, 108, 112, 147, 148, 161, 225–227, 239–259, 272, 274
- matrix function, 152
- matrix gradient, 191
- matrix inverse, 107
- matrix multiplication, 75–99, 528
 - Cayley, 75, 94
 - CUDA, 560
 - Hadamard, 94
 - inner product, 96–99
 - Kronecker, 94–96
 - MapReduce, 531
 - Strassen algorithm, 529–531
- matrix norm, 164–171
 - orthogonally invariant, 164
- matrix normal distribution, 218
- matrix of type 2, 58, 383
- matrix pencil, 161
- matrix polynomial, 78
 - Exercise 3.26*:, 181
- matrix random variable, 218–220
- matrix storage mode, 546–548
- Matrix Template Library, 569
- max norm, 28
- maximal linearly independent subset, 12
- maximum likelihood, 204–206
- MCMC (Markov chain Monte Carlo), 445
- mean, 35, 37
- mean vector, 35
- message passing, 557
- Message Passing Library, 557
- metric, 32, 175
- metric space, 32
- Microsoft R Open, 578
- MIL-STD-1753 standard, 478
- Minkowski inequality, 27
 - Exercise 2.11b*., 52
- Minkowski norm, 27
- minor, 67, 597
- missing data, 435–438, 571
 - representation of, 462, 473
- MKL (Intel Math Kernel Library), 555, 578
- mobile Jacobi scheme, 315
- modified Cholesky decomposition, 436
- “modified” Gauss-Newton, 203
- “modified” Gram-Schmidt (see also Gram-Schmidt transformation), 40
- Moore-Penrose inverse, 127–129, 248, 249, 291
 - relation to QR factorization, 249
- MPI (message passing interface), 557, 559
- MPL (Message Passing Library), 557
- multicollinearity, 265, 406
- multigrid method, 283
- multiple precision, 466, 491
- multiplicity of an eigenvalue, 144
- multivariate gamma function, 220
- multivariate linear regression, 418–422
- multivariate normal distribution, 217–219, 399, 441
 - singular, 217, 432
- multivariate random variable, 215–220
- $\mathcal{N}(\cdot)$, 126
- NA (“Not Available”), 462, 473, 571
- nabla (∇), 190, 191
- Nag Libraries, 556
- NaN (“Not-a-Number”), 473, 488
- NetCDF, 463
- netlib, xiv, 617
- network, 329–336, 391

- Newton's method, 198
- nilpotent matrix, 77, 174
- NMF (nonnegative matrix factorization), 257, 337
- noncentral chi-squared distribution, 400
 - PDF, equation (9.3), 400
- noncentral Wishart distribution
 - Exercise 4.12*, 222
- nonlinear regression, 201
- nonnegative definite matrix, 91, 159, 253, 344–350
 - summary of properties, 344–345
- nonnegative matrix, 258, 369
- nonnegative matrix factorization, 257, 337
- nonsingular matrix, 100, 111
- norm, 25–30
 - convexity, 26
 - equivalence of norms, 29, 33, 170
 - of a matrix, 164–171
 - orthogonally invariant, 164
 - of a vector, 27–31
 - weighted, 28, 93
- normal distribution, 217–219
 - matrix, 218
 - multivariate, 217–219
- normal equations, 256, 289, 404, 420
- normal matrix, 342
 - Exercise 8.1*, 394
 - circulant matrix, 384
- normal vector, 34
- normalized floating-point numbers, 468
- normalized generalized inverse (see also Moore-Penrose inverse), 128
- normalized vector, 31
- normed space, 25
- not-a-number (“NaN”), 473
- NP-complete problem, 504
- nuclear norm, 169
- null space, 126, 127, 144
- nullity, 126
- `numpy`, 556, 569
- Nvidia, 559

- $O(\cdot)$, 497, 503, 591
- $o(\cdot)$, 497, 591
- oblique projection, 356
- Octave (software), 578
- OLS (ordinary least squares), 288

- one vector, 16, 34
- online algorithm, 512
- online processing, 512
- open-source, 538
- OpenMP, 557, 559
- operator matrix, 80, 273
- operator norm, 165
- optimal design, 438–440
- optimization of vector/matrix functions, 196–212
 - constrained, 206–211
 - least squares, 200–204, 206–211
- order of a graph, 329
- order of a vector, 4
- order of a vector space, 15
- order of computations, 503
- order of convergence, 497
- order of error, 497
- ordinal relations among matrices, 92, 348
- ordinal relations among vectors, 16
- orthogonal array, 380
- orthogonal basis, 40–41
- orthogonal complement, 34, 126, 131
- orthogonal distance regression, 299–302, 405
- orthogonal group, 133, 220
- orthogonal matrices, binary relationship, 98
- orthogonal matrix, 131–134, 228
- orthogonal residuals, 299–302, 405
- orthogonal transformation, 228
- orthogonal vector spaces, 34, 131
- orthogonal vectors, 33
 - Exercise 2.6*, 52
- orthogonalization (Gram-Schmidt transformations), 38, 252, 524
- orthogonally diagonalizable, 147, 154, 338, 343, 423
- orthogonally invariant norm, 164, 167, 168
- orthogonally similar, 146, 154, 164, 168, 269, 343
- orthonormal vectors, 33
- out-of-core algorithm, 512
- outer product, 90, 245
 - Exercise 3.14*, 179
- outer product for matrix multiplication, 529

- outer pseudoinverse, 128, 129
- outer/inner products matrix, 357
- overdetermined linear system, 124, 255, 286
- overfitting, 298, 429
- overflow, in computer operations, 483, 487
- Overleaf, 539
- overloading, 11, 63, 165, 479, 489

- p -inverse (see also Moore-Penrose inverse), 128
- paging, 565
- parallel processing, 507, 508, 528, 530, 544, 557
- parallelogram equality, 27
- parallelotope, 74
- Parseval's identity, 41, 169
- partial ordering, 16, 92, 348
 - Exercise 8.2a*, 394
- partial pivoting, 275
- partitioned matrix, 61, 79, 131
 - determinant, 70, 122
 - sum of squares, 361, 413, 420
- partitioned matrix, inverse, 122, 131
- partitioning sum of squares, 361, 413, 420
- PBLAS (parallel BLAS), 558
- pencil, 161
- permutation, 66
- permutation matrix, 80, 86, 273, 377
- Perron root, 371, 374
- Perron theorem, 371
- Perron vector, 371, 375, 445
- Perron-Frobenius theorem, 374
- pivoting, 84, 244, 249, 275
- PLASMA, 559
- polar cone, 45
- polynomial in a matrix, 78
 - Exercise 3.26*., 181
- polynomial order, 503
- polynomial regression, 379
- polynomial, evaluation of, 512
- pooled variance-covariance matrix, 368
- population model, 445
- portability, 481, 494, 540
- positive definite matrix, 91, 101, 159–160, 253, 346–350, 422
 - summary of properties, 346–348
- positive matrix, 258, 369
- positive semidefinite matrix, 91
- positive stable, 159, 393
- power method for eigenvalues, 311–313
- precision, 472–480, 491
 - arbitrary, 466
 - double, 472, 480
 - extended, 472
 - half precision, 480
 - infinite, 466
 - multiple, 466, 491
 - single, 472, 480
- preconditioning, 282, 310, 525
 - for eigenvalue computations, 310
 - in the conjugate gradient method, 282
- primitive Markov chain, 445
- primitive matrix, 375, 445
- principal axis, 36
- principal components, 422–426
- principal components regression, 428
- principal diagonal, 56
- principal minor, 72, 104, 598
- principal submatrix, 62, 104, 241, 344, 347
 - leading, 62, 348
- probabilistic error bound, 497
- programming model, 463, 544
- projected gradient, 207
- projected Hessian, 207
- projection (of a vector), 20, 36
- projection matrix, 356–357, 408
- projective transformation, 228
- proper value (see also eigenvalue), 135
- PSBLAS (parallel sparse BLAS), 558
- pseudo-correlation matrix, 437
- pseudoinverse (see also Moore-Penrose inverse), 128
- PV-Wave (software), 6, 570
- Pythagorean theorem, 27
- Python, 478, 546, 569, 570

- Q-convergence, 509
- QL factorization, 247
- QR factorization, 246–252
 - and a generalized inverse, 249
 - computing, 558, 574
 - matrix rank, 250
 - skinny, 247

- QR method for eigenvalues, 316–318
- quadratic convergence, 509
- quadratic form, 91, 93
- quasi-Newton method, 199
- quotient space, 115
- R (software), 546, 570–578
 - Microsoft R Open, 578
 - Rcpp, 577
 - RcppArmadillo, 577
 - roxygen, 577
 - RPy, 578
 - RStudio, 578
 - Spotfire S+ (software), 578
- radix, 467
- random graph, 337
- random matrix, 218–220
 - BMvN distribution, 219
 - computer generation *Exercise 4.10.*, 221
 - correlation matrix, 441
 - Haar distribution, 219
 - Exercise 4.10.*, 221
 - normal, 218
 - orthogonal *Exercise 4.10.*, 221
 - rank *Exercise 4.11.*, 222
 - Wishart, 122, 346, 421, 432
 - Exercise 4.12.*, 222
- random number generation, 440–442
 - random matrices *Exercise 4.10.*, 221
- random variable, 215
- range of a matrix, 55
- rank deficiency, 100, 144
- rank determination, 532
- rank of a matrix, 99–122, 250, 431, 532
 - of idempotent matrix, 351
 - rank-revealing QR, 250, 431
 - statistical tests, 431–435
- rank of an array, 5
- rank reduction, 533
- rank(\cdot), 99
- rank, linear independence, 99, 532
- rank, number of dimensions, 5
- rank-one decomposition, 164
- rank-one update, 234, 285
- rank-revealing QR, 250, 431, 532
- rate constant, 509
- rate of convergence, 509
- rational fraction, 491
- Rayleigh quotient, 90, 157, 209, 392
- Rcpp, 577
- RcppBlaze, 559
- RCR (Replicated Computational Results), 552
- real numbers, 466
- real-time algorithm, 512
- recursion, 511
- reduced gradient, 207
- reduced Hessian, 207
- reduced rank regression problem, 432
- reducibility, 311, 334–336, 372
 - Markov chains, 444
- reflection, 231–233
- reflector, 233
- reflexive generalized inverse, 128
- register, in computer processor, 485
- regression, 401–422, 426–431
- regression variable selection, 427
- regression, nonlinear, 201
- regular graph, 329
- regular matrix (see also diagonalizable matrix), 149
- regularization, 298, 405, 429
- relative error, 484, 494, 526
- relative spacing, 470
- Reliable Computing*, 493
- Replicated Computational Results (RCR), 552
- Reproducible R Toolkit, 578
- reproducible research, 551–552, 578
- residue arithmetic, 493
- restarting, 525
- reverse communication, 564
- $\rho(\cdot)$ (spectral radius), 142
- Richardson extrapolation, 510
- ridge regression, 270, 362, 405, 418, 429
 - Exercise 9.11a.*, 453
- right direct product, 95
- right inverse, 108
- robustness (algorithm or software), 500
- root of a function, 487
- root-free Cholesky, 255
- Rosser test matrix, 550
- rotation, 229–232, 236
- rounding, 473
 - rounding error, 487, 495
- row echelon form, 111
- row rank, 99

- row space, 56
- row-major, 522, 539, 545
- row-sum norm, 166
- roxygen, 577
- RPy, 578
- RQ factorization, 247
- RStudio, 578

- S (software), 570
- Samelson inverse, 31
- sample variance, computing, 501
- saxpy**, 12
- scalability, 506
- ScaLAPACK, 558
- scalar, 11
- scalar product, 23
- scaled matrix, 365
- scaled vector, 49
- scaling of a vector or matrix, 269
- scaling of an algorithm, 503, 506
- Schatten p norm, 169
- Schur complement, 121, 412, 421
- Schur factorization, 147–148
- Schur norm (see also Frobenius norm), 167
- SDP (semidefinite programming), 346
- Seidel adjacency matrix, 332
- self-adjoint matrix (see also Hermitian matrix), 56
- semidefinite programming (SDP), 346
- seminorm, 25
- semisimple eigenvalue, 144, 149
- sequences of matrices, 171
- sequences of vectors, 32
- shape of matrix, 5
- shearing transformation, 228
- Sherman-Morrison formula, 285, 415
- shifting eigenvalues, 310
- shrinkage, 405
- side effect, 554
- $\sigma(\cdot)$ (sign of permutation), 66, 86
- $\sigma(\cdot)$ (spectrum of matrix), 141
- sign bit, 465
- sign(\cdot), 16
- significand, 467
- similar canonical form, 149
- similar matrices, 146
- similarity matrix, 369
- similarity transformation, 146–148, 313, 317
- simple eigenvalue, 144
- simple graph, 329
- simple matrix (see also diagonalizable matrix), 149
- single precision, 472, 480
- singular matrix, 100
- singular multivariate normal distribution, 217, 432
- singular value, 162, 425, 532
 - relation to eigenvalue, 163
- singular value decomposition, 161–164, 320, 336, 425, 532
 - uniqueness, 163
- skew diagonal element, 57
- skew diagonal matrix, 57
- skew symmetric matrix, 56, 60
- skew upper triangular matrix, 58, 388
- skinny QR factorization, 247
- smoothing matrix, 361, 418
- software testing, 548–551
- SOR (method), 279
- span(\cdot), 14, 21, 55
- spanning set, 14, 21
 - of a cone, 44
- Spark (software system), 544
- sparse matrix, 59, 259, 277, 523, 526, 556, 557
 - index-index-value, 548
 - software, 557
 - CUDA, 560
 - storage mode, 548
- spectral circle, 142
- spectral condition number, 268, 269, 290
- spectral decomposition, 155, 163
- spectral norm, 166, 169
- spectral projector, 155
- spectral radius, 142, 166, 170, 278
- spectrum of a graph, 392
- spectrum of a matrix, 141–145
- splitting extrapolation, 511
- Spotfire S+ (software), 578
- square root matrix, 160, 252, 254, 345
- stability, 276, 500
- standard deviation, 49, 364
 - computing the standard deviation, 501

- Standard Template Library, 569
- standards (see also specific standard), 460
- stationary point of vector/matrix functions, 198
- statistical reference datasets (StRD), 551
- statlib**, xiv, 617
- steepest descent, 198, 199
- Stiefel manifold, 133
- stiff data, 502
- stochastic matrix, 377
- stochastic process, 442–450
- stopping criterion, 508
- storage mode, for matrices, 546–548
- storage unit, 464, 467, 480
- Strassen algorithm, 529–531
- StRD (statistical reference datasets), 551
- stride, 522, 539, 554
- string, character, 461
- strongly connected graph, 334
- submatrix, 61, 79
- subspace of vector space, 17
- successive overrelaxation, 279
- summation, 485, 486
- summing vector, 34
- Sun ONE Studio Fortran 95, 493
- superlinear convergence, 509
- SVD (singular value decomposition), 161–164, 320, 336, 425, 532
 - uniqueness, 163
- sweep operator, 413
- Sylvester’s law of nullity, 117
- symmetric matrix, 56, 60, 112, 153–160, 338–343
 - eigenvalues/vectors, 153–160
 - equivalent forms, 112
 - inverse of, 120
 - summary of properties, 338
- symmetric pair, 319
- symmetric storage mode, 61, 547

- Taylor series, 188, 198
- Template Numerical Toolkit, 569
- tensor, 5
- term-document matrix, 336
- test problems for algorithms or software, 527, 548–551
 - Exercise 3.24*:, 180
 - consistency test, 527, 551
 - Ericksen matrix, 549
 - Exercise 12.8*:, 582
 - Hilbert matrix, 548
 - Matrix Market, 550
 - randomly generated data, 549
 - Exercise 11.5*:, 536
 - Rosser matrix, 550
 - StRD (statistical reference datasets), 551
 - Wilkinson matrix, 550
 - Exercise 12.9*:, 582
 - Wilkinson’s polynomial, 499
- testable hypothesis, 410
- testing software, 548–551
- thread, 544
- Tikhonov regularization, 299, 429
- time series, 447–450
 - variance-covariance matrix, 383, 449
- Toeplitz matrix, 382, 449
 - circulant matrix, 383
 - inverse of, 383
 - Exercise 8.12*:, 395
 - Exercise 12.12*:, 583
- total least squares, 299, 405
- tr(\cdot), 65
- trace, 65
 - derivative of, 195
 - of Cayley product, 87
 - of idempotent matrix, 351
 - of inner product, 92
 - of Kronecker product, 96
 - of matrix inner product, 97
 - of outer product, 92
 - relation to eigenvalues, 141
- trace norm, 169
- transition matrix, 443
- translation transformation, 232
- transpose, 59
 - determinant of, 70
 - generalized inverse of, 124
 - inverse of, 107
 - norm of, 164
 - of Cayley product of matrices, 76
 - of Kronecker product, 95
 - of partitioned matrices, 63
 - of sum of matrices, 63
 - trace of, 65

- trapezoidal matrix, 58, 241, 246
- triangle inequality, 25, 164
 - Exercise 2.11b*, 52
- triangular matrix, 57, 84, 240
 - determinant of, 70
 - inverse of, 121
 - multiplication, 78
- tridiagonal matrix, 58
- triple scalar product *Exercise 2.19c*, 54
- triple vector product *Exercise 2.19d*, 54
- truncation error, 42, 99, 498
- twos-complement representation, 465, 483
- type 2 matrix, 58, 383

- ulp (“unit in the last place”), 471
- underdetermined linear system, 123
- underflow, in computer operations, 470, 487
- Unicode, 461
- union of vector spaces, 18
- unit in the last place (ulp), 471
- unit roundoff, 470
- unit vector, 16, 22, 36, 76
- unitarily diagonalizable, 147, 343, 387
- unitarily similar, 146
- unitary matrix, 132
- unrolling do-loop, 565
- updating a solution, 285, 293, 415–417
 - regression computations, 415–417
- upper Hessenberg form, 59, 316
- upper triangular matrix, 57
- usual norm (see also Frobenius norm), 167

- $V(\cdot)$ (variance operator), 49, 216
- $\mathcal{V}(\cdot)$ (vector space), 21, 55
- Vandermonde matrix, 379
 - Fourier matrix, 384
- variable metric method, 199
- variable selection, 427
- variance, computing, 501
- variance-covariance matrix, 216, 219
 - Kronecker structure, 219, 419
 - positive definite approximation, 436
 - sample, 365, 422
- $\text{vec}(\cdot)$, 61
- vec-permutation matrix, 82
- $\text{vecdiag}(\cdot)$, 56

- $\text{vech}(\cdot)$, 61
- vector, 4
 - centered vector, 48
 - mean vector, 35
 - normal vector, 34
 - normalized vector, 31
 - null vector, 15
 - one vector, 15, 34
 - “row vector”, 89
 - scaled vector, 49
 - sign vector, 16
 - summing vector, 15, 34
 - unit vector, 16
 - zero vector, 15
- vector derivative, 183–220
- vector processing, 557
- vector space, 13–15, 17–23, 55, 64, 126, 127
 - basis, 21–23
 - definition, 13
 - dimension, 14
 - direct product, 20
 - direct sum, 18–20
 - direct sum decomposition, 19
 - essentially disjoint, 15
 - intersection, 18
 - null vector space, 13
 - of matrices, 64
 - order, 15
 - set operations, 17
 - subspace, 17
 - union, 18
- vector subspace, 17
- vectorized processor, 508
- vertex of a graph, 329
- volume as a determinant, 73, 217

- weighted graph, 329
- weighted least squares, 414
 - with equality constraints *Exercise 9.4d*, 451
- weighted norm, 28, 93
- Wilk’s Λ , 421
- Wilkinson matrix, 550
- Wishart distribution, 122, 346
 - Exercise 4.12*, 222
- Woodbury formula, 285, 415
- word, computer, 464, 467, 480

XDR (external data representation),
481

Yule-Walker equations, 449

Z-matrix, 393

zero matrix, 76, 99

zero of a function, 487

zero vector, 15