
Index

- absolute error, [541](#), [552](#), [584](#)
- ACM Transactions on Mathematical Software*, [611](#)
- $\text{adj}(\cdot)$, [90](#)
- adjacency matrix, [363–366](#), [422](#)
 - Exercise 8.20.*, [427](#)
 - augmented, [422](#)
- adjoint (see also conjugate transpose), [73](#)
- adjoint, classical (see also adjugate), [90](#)
- adjugate, [90](#), [650](#)
- adjugate and inverse, [136](#)
- affine group, [130](#), [204](#)
- affine space, [51](#)
- affine transformation, [214](#)
- Aitken’s integral, [340](#)
- $\mathcal{AL}(\cdot)$ (affine group), [130](#)
- algebraic multiplicity, [163](#)
- algorithm, [254](#), [556–570](#)
 - batch, [568](#)
 - definition, [566](#)
 - direct, [254](#)
 - divide and conquer, [563](#)
 - greedy, [563](#)
 - iterative, [254](#), [565–568](#), [623](#)
 - reverse communication, [623](#)
 - online, [568](#)
 - out-of-core, [568](#)
 - real-time, [568](#)
- algorithmically singular, [139](#)
- Anaconda, [612](#), [615](#), [628](#)
- $\text{angle}(\cdot, \cdot)$, [45](#)
- angle between matrices, [188](#)
- angle between vectors, [45](#), [215](#), [389](#)
- ANSI (standards), [533](#), [621](#), [622](#)
- antisymmetric matrix, [70](#)
- Apache Software Foundation, [518](#), [522](#)
- approximation and estimation, [469](#)
- approximation of a matrix, [194](#), [243](#), [371](#), [474](#), [590](#)
- approximation of a vector, [49–51](#)
- arithmetic mean, [43](#), [45](#)
- Arnoldi method, [310](#)
- artificial ill-conditioning, [259](#)
- ASCII code, [519](#)
- association matrix, [360–370](#), [389](#), [397](#), [401–403](#)
 - adjacency matrix, [363–364](#), [422](#)
 - connectivity matrix, [363–364](#), [422](#)
 - dissimilarity matrix, [401](#)
 - distance matrix, [401](#)
 - incidence matrix, [363–364](#), [422](#)
 - similarity matrix, [401](#)
- ATLAS (Automatically Tuned Linear Algebra Software), [614](#)
- augmented adjacency matrix, [422](#)
- augmented connectivity matrix, [422](#)
- Automatically Tuned Linear Algebra Software (ATLAS), [614](#)
- autoregressive process, [481–499](#)
- axpy, [18](#), [59](#), [101](#), [613](#), [614](#)
- axpy elementary operator matrix, [101](#)
- back substitution, [264](#), [439](#)
- backward error analysis, [552](#), [558](#)
- Banach space, [41](#)
- Banachiewicz factorization, [241](#)

- banded matrix, 72
 - inverse, 139
- Bartlett decomposition, 378
- base, 526
- base point, 525
- basis, 29–30
 - Exercise 2.19.*, 65
 - orthonormal, 48–49
- batch algorithm, 568
- Bauer-Fike theorem, 299
- Beowulf (cluster computing), 618
- bias, in exponent of floating-point number, 526
- big endian, 538
- big integer, 524, 549
- big O (order), 554, 560, 643
- big omega (order), 555
- bilinear form, 108, 152
- bit, 519
- bitmap, 520
- BLACS (software), 616, 617
- BLAS (software), 612–615
 - CUDA, 619
 - PBLAS, 616, 617
 - PLASMA, 617
 - PSBLAS, 616
- block diagonal matrix, 77
 - determinant of, 83
 - inverse of, 139
 - multiplication, 95
- BMvN distribution, 343, 607
- Bolzano-Weierstrass theorem for orthogonal matrices, 151
- Boolean matrix, 422
- Box M statistic, 400
- bra-ket notation, 31
- byte, 519
- C (programming language), 533, 548, 627–628
- C++ (programming language), 534, 627–628
- cancellation error, 544, 558
- canonical form, equivalent, 130
- canonical form, similar, 167
- canonical singular value factorization, 179
- Cartesian geometry, 43, 86
- catastrophic cancellation, 544
- Cauchy matrix, 421
- Cauchy-Schwarz inequality, 32, 116
- Cauchy-Schwarz inequality for matrices, 116, 204
- Cayley multiplication, 91, 111
- Cayley-Hamilton theorem, 157
- CDF (Common Data Format), 521
- centered matrix, 277, 395
- centered vector, 58
- chaining of operations, 542
- change-of-variables method, 339
- character data, 520
- character string, 520
- characteristic equation, 156
- characteristic function, 335
- characteristic polynomial, 156
- characteristic value (see also eigenvalue), 153
- characteristic vector (see also eigenvector), 153
- chasing, 309
- Chebyshev norm, 35
- chi-squared distribution, 346
 - noncentral, 346
 - PDF, equation (7.53), 346
- Cholesky decomposition, 240–242, 377, 474
 - computing, 617
 - root-free, 241
- circulant matrix, 415
- classification, 421
- cluster analysis, 421
- cluster computing, 618
- coarray (Fortran construct), 622
- Cochran's theorem, 347, 385–387
- cofactor, 89, 650
- collinearity, 255, 435, 468
- column rank, 117
- column space, 69, 96, 122, 123
- column-major, 580, 598, 604
- column-sum norm, 185
- Common Data Format (CDF), 521
- companion matrix, 157, 297
- compatible linear systems, 124
- compensated summation, 543
- complementary projection matrix, 388
- complementary vector spaces, 27
- complete graph, 361
- complete pivoting, 265

- complete space, 41
- completing the Gramian, 196
- complex numbers, vectors, and matrices, 41, 150, 418
- computer algebra, 601
- Conda, 612
- condition (problem or data), 556
- condition number, 255, 260, 279, 463, 464, 557, 559, 581, 591
 - computing the number, 591
 - inverse of matrix, 256, 260
 - nonfull rank matrices, 279
 - nonsquare matrices, 279
 - sample standard deviation, 559
- conditional inverse, 146
- cone, 51–54, 359
 - Exercise 2.20*., 65
 - convex cone, 52, 378
 - of nonnegative definite matrices, 378
 - of nonnegative matrices, 403
 - of positive definite matrices, 381
 - of positive matrices, 403
- conference matrix, 413
- configuration matrix, 401
- conjugate gradient method, 269–273
 - preconditioning, 271
- conjugate norm, 111
- conjugate transpose, 73, 150
- conjugate vectors, 110, 152
- connected vertices, 361, 366
- connectivity matrix, 363–366, 422
 - Exercise 8.20*., 427
 - augmented, 422
- consistency property of matrix norms, 183
- consistency test, 584, 609
- consistent system of equations, 123, 262, 267
- constrained least squares, equality constraints, 443
 - Exercise 9.3d*., 507
- continuous function, 320
- contrast, 454
- convergence criterion, 566
- convergence of a sequence of matrices, 151, 171, 191
- convergence of a sequence of vectors, 40
- convergence of powers of a matrix, 191, 407
- convergence rate, 566
- convex combination, 18
- convex cone, 52–54, 378, 381, 403
- convex function, 33, 484
- convex optimization, 378
- convex set, 18
- convexity, 33
- coordinate, 3
- Cor($\cdot, \cdot$), 60
- correlation, 60, 97
- correlation matrix, 397, 459
 - Exercise 7.12*., 356
 - positive definite approximation, 474
 - pseudo-correlation matrix, 475
 - sample, 460
- cost matrix, 401
- Cov($\cdot, \cdot$), 59
- covariance, 59
- covariance matrix, *see* variance-covariance matrix
- CRAN (Comprehensive R Archive Network), 10, 611, 631
- cross product of vectors, 55
 - Exercise 2.21*., 66
- cross products matrix, 242, 390
- cross products, computing sum of
 - Exercise 10.17c*., 577
- Crout method, 231
- cuBLAS, 619
- CUDA, 618
- curl, 327
- curse of dimensionality, 568
- cuSPARSE, 619
- D-optimality, 591
- data frame, 501
- daxpy, 18
- decomposable matrix, 405
- decomposition, *see also* factorization of a matrix
 - additive, 385, 386
 - Bartlett decomposition, 378
 - Cholesky, 240
 - multiplicative, 126
 - nonnegative matrix factorization, 244, 369
 - singular value decomposition, 178–181, 312, 368, 462, 590
 - spectral decomposition, 173

- defective (deficient) matrix, 167, 169
- deficient (defective) matrix, 167, 169
- deflation, 300–302
- degrees of freedom, 392, 394, 441, 468
- del (∇), 326
- derivative with respect to a matrix, 328–330
- derivative with respect to a vector, 323–328
- $\det(\cdot)$, 81
- determinant, 81–87
 - computing, 590
 - derivative of, 330
 - Jacobian, 340
 - of block diagonal matrix, 83
 - of Cayley product, 106
 - of diagonal matrix, 83
 - of elementary operator matrix, 104
 - of inverse, 135
 - of Kronecker product, 113
 - of nonnegative definite matrix, 377
 - of partitioned matrix, 83, 140
 - of permutation matrix, 105
 - of positive definite matrix, 379
 - of transpose, 82
 - of triangular matrix, 83
 - relation to eigenvalues, 159
 - relation to geometric volume, 86, 340
- $\text{diag}(\cdot)$, 74, 75, 648
 - with matrix arguments, 77
- diagonal element, 70
- diagonal expansion, 86
- diagonal factorization, 167, 171
- diagonal matrix, 71
 - determinant of, 83
 - inverse of, 138
 - multiplication, 95
- diagonalizable matrix, 167–171, 298, 375
 - orthogonally, 173
 - singular value decomposition, 181
 - unitarily, 165, 375, 419
- diagonally dominant matrix, 71, 77, 119, 379
- differential, 322
- differentiation of vectors and matrices, 317–330
- digraph, 364
 - of a matrix, 365
- $\dim(\cdot)$, 21
- dimension of matrix, 4
- dimension of vector space, 21
- dimension reduction, 28, 388, 463
- direct method for solving linear systems, 262–266
- direct product (of matrices), 112
- direct product (of sets), 3
- direct product (of vector spaces), 28
 - basis for, 30
- direct sum decomposition of a vector space, 27
- direct sum of matrices, 78
- direct sum of vector spaces, 25–27, 79
 - basis for, 30
 - direct sum decomposition, 27
- directed dissimilarity matrix, 401
- direction cosines, 46, 217
- discrete Fourier transform, 416
- discrete Legendre polynomials, 412
- discretization error, 555, 567
- dissimilarity matrix, 401
- distance, 39
 - between matrices, 195
 - between vectors, 40
- distance matrix, 401
- distributed computing, 522, 565, 603
- distributed linear algebra machine, 617
- distribution vector, 409
- $\text{div}(\cdot)$, 326
- divergence, 326
- divide and conquer, 563
- document-term matrix, 368
- dominant eigenvalue, 160
- Doolittle method, 231
- dot product of matrices, 115
- dot product of vectors, 31, 108
- double cone, 51
- double precision, 531, 537
- doubly stochastic matrix, 409
- `dplyr` (R package), 502
- dual cone, 52
- E_{pq} , $E_{(\pi)}$, $E_p(a)$, $E_{pq}(a)$ (elementary operator matrices), 103
- $E(\cdot)$ (expectation operator), 335
- echelon form, 131
- edge of a graph, 361
- EDP (exact dot product), 551

- effective degrees of freedom, 394, 468
- efficiency, computational, 560–565
- eigenpair, 152
- eigenspace, 162
- eigenvalue, 152–181, 186, 297–314
 - computing, 298–310, 617
 - Jacobi method, 305–308
 - Krylov methods, 310
 - power method, 303–305
 - QR method, 308–310
 - of a graph, 423
 - of a polynomial *Exercise 3.26*:, 205
 - relation to singular value, 180
 - upper bound on, 160, 163, 298
- eigenvector, 152–181, 297–314
 - left eigenvector, 155, 176
- eigenvectors, linear independence of, 161
- EISPACK, 615
- elastic net regression, 468
- elementary operation, 98
- elementary operator matrix, 98–105, 119, 228, 263
 - eigenvalues, 154
- elliptic metric, 111
- elliptic norm, 111
- endian, 538
- equivalence of norms, 37, 40, 189
- equivalence relation, 478
- equivalent canonical factorization, 132
- equivalent canonical form, 130, 132
- equivalent matrices, 117, 130
- error bound, 554
- error in computations
 - cancellation, 544, 558
 - error-free computations, 550
 - measures of, 541, 552–554, 584
 - rounding, 544, 552, 553
 - Exercise 10.9*:, 575
- error of approximation, 555
 - discretization, 555
 - truncation, 556
- error, measures of, 262, 541, 552–554, 584
- error-free computations, 550
- errors-in-variables, 435
- essentially disjoint vector spaces, 24, 79
- estimable combinations of parameters, 453
- estimation and approximation, 469
- Euclidean distance, 40, 401
- Euclidean distance matrix, 401
- Euclidean matrix norm (see also Frobenius norm), 187
- Euclidean vector norm, 35
- Euler’s constant *Exercise 10.2*:, 574
- Euler’s integral, 645
- Euler’s rotation theorem, 217
- exact computations, 550
- exact dot product (EDP), 551
- exception, in computer operations, 540, 545
- expectation, 335–343
- exponent, 526
- exponential order, 560
- exponential, matrix, 171, 318
- extended precision, 531
- extrapolation, 567
- factorization of a matrix, 93, 126, 132, 165, 167, 178, 181, 211–213, 226–245, 262, 264
 - Banachiewicz factorization, 241
 - Bartlett decomposition, 378
 - canonical singular value factorization, 179
 - Cholesky factorization, 240–242
 - diagonal factorization, 167
 - equivalent canonical factorization, 132
 - full rank factorization, 126
 - Gaussian elimination, 262
 - LQ factorization, 233
 - LU or LDU factorization, 226–232
 - nonnegative matrix factorization, 244, 369
 - orthogonally diagonal factorization, 165, 181
 - QL factorization, 233
 - QR factorization, 232–238
 - root-free Cholesky, 241
 - RQ factorization, 233
 - Schur factorization, 166
 - singular value factorization, 178–181, 312, 368, 462, 590
 - square root factorization, 183
 - unitarily diagonal factorization, 165, 181

- fan-in algorithm, 543, 563
- fast Fourier transform (FFT), 418
- fast Givens rotation, 225, 583
- fill-in, 245, 584
- Fisher information, 492
- fixed-point representation, 524
- flat, 51
- floating-point representation, 525
- FLOP, or flop, 562
- FLOPS, or flops, 562
- Fortran, 534–536, 562, 605, 625–627
- Fourier coefficient, 49, 50, 117, 175, 181, 188
- Fourier expansion, 44, 49, 116, 175, 181, 188
- Fourier matrix, 416
- Free Software Foundation, 518
- Frobenius norm, 187–188, 191, 195, 306, 372, 401
 - Frobenius p norm, 189
- full precision, 537
- full rank, 118, 119, 122, 128
- full rank factorization, 126
- full rank partitioning, 122, 140
- g_{12} inverse, 146, 437
- g_4 inverse (see also Moore–Penrose inverse), 147
- gamma function, 343, 645
- GAMS (*Guide to Available Mathematical Software*), 598
- Gauss–Markov theorem, 455
- Gauss–Newton method, 490
- Gauss–Seidel method, 267
- Gaussian elimination, 102, 262, 309
- Gaussian matrix, 102, 228
- gemm (general matrix-matrix), 615
- gemv (general matrix-vector), 614
- general linear group, 130, 151
- generalized eigenvalue, 177, 311
- generalized inverse, 142–149, 235, 391, 437
 - relation to QR factorization, 236
- generalized least squares, 443
- generalized least squares with equality constraints *Exercise 9.3d*, 507
- generalized linear model, 503
- generalized variance, 398
- generating set, 21, 29
 - of a cone, 52
- generation of random numbers, 11–12, 348–352
- geometric multiplicity, 162
- geometry, 43, 86, 213, 217
- Gershgorin disks, 163
- GitHub, 10, 597
 - Exercise 12.3*:, 633
- Givens transformation (rotation), 222–225, 309
 - QR factorization, 238
- $GL(\cdot)$ (general linear group), 130
- GMP (software library), 524, 549
 - Exercise 10.25*:, 578
 - Exercise 10.26*:, 578
- GMRES, 271
- GNU Scientific Library (GSL), 615
- GPU (graphical processing unit), 618
- graceful underflow, 528
- gradient, 324
 - projected gradient, 494
 - reduced gradient, 494
- gradient descent, 484, 486
- gradient of a function, 324, 325
- gradual underflow, 528, 545
- Gram–Schmidt transformation, 47, 48, 582
 - linear least squares, 279
 - QR factorization, 238
- Gramian matrix, 132, 134, 158, 179, 182, 242, 278, 389–391
 - completing the Gramian, 196
 - eigenvalues, 158
- graph of a matrix, 364
- graph theory, 7, 361–368, 421
- graphical processing unit (GPU), 618
- greedy algorithm, 563
- group, 130, 151
- GSL (GNU Scientific Library), 615
- guard digit, 542
- Haar distribution, 343, 607
 - Exercise 7.11*:, 355
 - Exercise 7.12*:, 356
- Haar invariant measure, 343
- Hadamard matrix, 412
- Hadamard multiplication, 111
- Hadamard’s inequality *Exercise 4.4*:, 9, 248

- Hadoop, 518, 522, 603
- Hadoop Distributed File System (HDFS), 522, 571
- half precision, 537
- Hankel matrix, 420
- Hankel norm, 421
- hat matrix, 392, 442
- HDF, HDF5 (Hierarchical Data Format), 521
- HDFS (Hadoop Distributed File System), 522, 571
- Helmert matrix, 410, 454
- Hemes formula, 275, 445
- Hermite form, 131
- Hermitian matrix, 70, 74
- Hessenberg matrix, 73, 308
- Hessian matrix, 328
 - projected Hessian, 494
 - reduced Hessian, 494
- Hessian of a function, 328
- hidden bit, 526
- Hierarchical Data Format (HDF), 521
- high-performance computing, 564
- Hilbert matrix, 607
- Hilbert space, 41, 187
- Hilbert-Schmidt norm (see also Frobenius norm), 187
- Hoffman-Wielandt theorem, 371
- Hölder norm, 35
- Hölder's inequality *Exercise 2.12a*, 64
- hollow matrix, 71, 401
- homogeneous coordinates, 218
 - in graphics applications, *Exercise 4.10*, 249
- homogeneous system of equations, 51, 141
- Horner's method, 570
- Householder transformation (reflection), 219–222, 237, 309
- hyperplane, 51
- hypothesis testing, 449
- idempotent matrix, 94, 130, 382–389
- identity matrix, 93
 - additive identity, 78
- IDL (software), 4
- IEC standards, 523
- IEEE standards, 523, 544
 - Standard 754, 531, 537, 544, 551
 - Standard P1788, 550
- IFIP Working Group 2.5, 518, 551
- ill-conditioned (problem or data), 254, 464, 556, 581
 - artificial, 259
 - stiff data, 559
- ill-posed problem, 139
- image data, 520
- IMSL Libraries, 615, 619–620
- incidence matrix, 363–366, 422
- incomplete data, 473–476
- incomplete factorization, 245, 584
- independence, linear, *see* linear independence
- independent vertices, 422
- index of a matrix, 130
- index of idempotent matrix, 94
- index of nilpotent matrix, 94
- index-index-value (sparse matrices), 607
- induced matrix norm, 184
- Inf, 12
- infinity, floating-point representation, 531, 532, 545
- infix operator, 546
- inner product, 30, 231
 - inner product of matrices, 115–117
 - inner product space, 32
- inner pseudoinverse, 146
- integer representation, 524
- integration, 331–333
- integration of vectors and matrices, 331
- Intel Math Kernel Library (MKL), 614, 633
- intersection graph, 366
- intersection of vector spaces, 25
- interval arithmetic, 549, 550
- invariance property, 213
- invariant distribution, 479
- invariant vector (eigenvector), 153
- inverse CDF method, 339
- inverse of a matrix, 125
 - determinant of, 135
 - generalized inverse, 142–149
 - Moore-Penrose inverse, 146–148
 - pseudoinverse, 147
 - Kronecker product, 136
 - left inverse, 126
 - Moore-Penrose inverse, 146–148
 - partitioned matrix, 140

- products or sums of matrices, 136
- pseudoinverse, 147
- right inverse, 126
- transpose, 125
- triangular matrix, 139
- inverse of a vector, 39
- IRLS (iteratively reweighted least squares), 287
- irreducible Markov chain, 478
- irreducible matrix, 303, 367–368, 404–409, 478
- `is.na`, 532
- `is.nan`, `is.nan`, 532
- ISO (standards), 533, 621, 622
- isometric matrix, 186
- isometric transformation, 214
- isotropic transformation, 214
- iterative method, 267, 273, 297, 565–568, 583
 - for solving linear systems, 267–273
 - iterative refinement, 273
- iteratively reweighted least squares, 287
- Jacobi method for eigenvalues, 305–308
- Jacobi transformation (rotation), 222
- Jacobian, 325, 340
- Jordan decomposition, 169
- Jordan form, 131
- Kalman filter, 559
- Kantorovich inequality, 382
- Karush-Kuhn-Tucker conditions, 498
- kind (for data types), 534
- Kronecker multiplication, 112–114
 - inverse, 136
 - properties, 113
 - symmetric matrices, 113, 174
 - diagonalization, 174
- Kronecker structure, 114, 343, 458
- Kronecker sum, 114
- Krylov method, 271, 310
- Krylov space, 271
- Kuhn-Tucker conditions, 498
- Kulisch accumulator, 551
- Kullback-Leibler divergence, 195
- L_1 , L_2 , and L_∞ norms
 - of a matrix, 185
 - of a symmetric matrix, 187
 - of a vector, 35
 - relations among, 190
- L_2 norm of a matrix (see also spectral norm), 186
- Lagrange multiplier, 443, 495
 - Exercise 9.3a*., 506
- Lagrangian function, 495
- Lanczos method, 310
- LAPACK, 266, 591, 615, 617
 - LAPACK95, 615
- Laplace expansion, 89
- Laplace operator (∇^2), 651
- Laplace operator (∇^2), 326
- Laplacian matrix, 423
- lasso regression, 468
- latent root (see also eigenvalue), 153
- LAV (least absolute values), 285
- LDU factorization, 226–232
- leading principal submatrix, 77, 380
- least absolute values, 285
- least squares, 242, 276–284, 487–491
 - constrained, 493–498
 - maximum likelihood, 452
 - nonlinear, 489–491
 - nonnegative, 289, 497
- least squares regression, 487
- left eigenvector, 155, 176
- left inverse, 126
- length of a vector, 2, 35, 38
- Leslie matrix, 410, 480
 - Exercise 8.10*., 426
 - Exercise 9.15*., 509
- Levenberg-Marquardt method, 491
- leverage, 442
 - Exercise 9.18*., 510
- life table, 480
- likelihood function, 491
- line, 51
- linear convergence, 566
- linear estimator, 453
- linear independence, 18, 117
- linear independence of eigenvectors, 161
- linear programming, 378
- linear regression, 430–459, 463–468
 - variable selection, 464
- LINPACK, 266, 591, 615
- Linux Foundation, 518
- little endian, 538
- little o (order), 555, 643

- little omega (order), 555
- log order, 560
- log-likelihood function, 492
- Longley data *Exercise 9.19*, 511
- loop unrolling, 624
- Lorentz cone, 52
- lower triangular matrix, 71
- L_p norm
 - of a matrix, 185
 - of a vector, 35–36, 320
- LQ factorization, 233
- LR method, 298
- LU factorization, 226–232
 - computing, 617
- M-matrix, 425
- MACHAR, 536
 - Exercise 10.3(d)i*, 574
- machine epsilon, 12, 528
- Mahalanobis distance, 111, 397
- Manhattan norm, 35
- manifold of a matrix, 69
- Maple (software), 621
- MapReduce, 522, 570, 588, 603
- Markov chain, 476–479
- Markov chain Monte Carlo (MCMC), 479
- Mathematica (software), 621
- Matlab (software), 605, 628–630
- `matlib` (R package), 197
- matrix, 3
- Matrix** (R package), 197
- matrix derivative, 317–330
- matrix exponential, 171, 318
- matrix factorization, 93, 126, 132, 165, 167, 178, 181, 211–213, 226–245, 262, 264
- matrix function, 171
- matrix gradient, 325
- matrix inverse, 125
- matrix multiplication, 91–117, 585
 - Cayley, 91, 111
 - CUDA, 619
 - Hadamard, 111
 - inner product, 115–117
 - Kronecker, 112–114
 - MapReduce, 588
 - Strassen algorithm, 587–588
- matrix norm, 183–191
 - orthogonally invariant, 184
- matrix normal distribution, 342
- matrix of type 2, 72, 415
- matrix pencil, 178
- matrix polynomial, 95
 - Exercise 3.26*:, 205
- matrix random variable, 337, 342–343
- matrix storage mode, 605–607
- Matrix Template Library, 627
- max norm, 35
- maximal linearly independent subset, 19
- maximum likelihood, 452, 491–493, 503
 - least squares, 452
- MCMC (Markov chain Monte Carlo), 479
- mean, 43, 45
- mean vector, 42
- message passing, 616
- Message Passing Library, 616
- metric, 39, 195
- metric space, 40
- Microsoft R Open, 633
- MIL-STD-1753 standard, 536
- Minkowski inequality, 35
 - Exercise 2.12b*:, 64
- Minkowski norm, 35
- minor, 87, 649
- missing data, 473–476, 500
 - casewise deletion, 474, 500
 - pairwise deletion, 474, 501
 - representation of, 520, 532
- MKL (Intel Math Kernel Library), 614, 633
- mobile Jacobi scheme, 307
- modified Cholesky decomposition, 474
- “modified” Gauss-Newton, 490
- “modified” Gram-Schmidt (see also Gram-Schmidt transformation), 48
- moment-generating function (MGF), 335
- Moore-Penrose inverse, 146–148, 235, 236, 281, 437
 - relation to QR factorization, 236
- MPI (message passing interface), 616, 618
- MPL (Message Passing Library), 616
- multicollinearity, 255, 435

- multigrid method, 273
- multiple precision, 524, 549
- multiplicity of an eigenvalue, 163
- multivariate gamma function, 343
- multivariate linear regression, 448–449, 457–459
- multivariate normal distribution, 339–343, 349
 - singular, 340, 470
- multivariate random variable, 334–343
- $\mathcal{N}(\cdot)$, 144
- NA (“Not Applicable”), 12, 500, 520, 532
- nabla (∇), 324, 325
- Nag Libraries, 615
- NaN (“Not-a-Number”), 12, 532, 545
- NetCDF, 521
- netlib, xvi
- network, 361–368, 423
- Newton’s method, 485
- nilpotent matrix, 94, 193
 - unipotent matrix, 95
- NMF (nonnegative matrix factorization), 244, 369
- noncentral chi-squared distribution, 346
 - PDF, equation (7.53), 346
- noncentral Wishart distribution
 - Exercise 7.9*, 354
- nonlinear regression, 487
- nonnegative definite matrix, 108, 181, 240, 376–382
 - summary of properties, 376–377
- nonnegative least squares, 289, 497
- nonnegative matrix, 244, 401
- nonnegative matrix factorization, 244, 369
- nonsingular matrix, 119, 128
- norm, 33–38
 - convexity, 33
 - equivalence of norms, 37, 40, 189
 - of a matrix, 183–191
 - orthogonally invariant, 184
 - of a vector, 35–39
 - weighted, 36, 111
- normal distribution, 339–343
 - matrix, 342
 - multivariate, 339–343
- normal equations, 242, 278, 433, 449
- normal matrix, 374
 - Exercise 8.1*, 425
 - circulant matrix, 416
- normal vector, 42
- normalized floating-point numbers, 526
- normalized generalized inverse (see also Moore–Penrose inverse), 147
- normalized vector, 39
- normed space, 33
- not-a-number (“NaN”), 532
- NP-complete problem, 561
- nuclear norm, 189
- null space, 144, 145, 162
- nullity, 144
- numpy, 615, 628
- Nvidia, 618
- $O(\cdot)$, 554, 560, 643
- $o(\cdot)$, 555, 643
- oblique projection, 388
- Octave (software), 628–630
- OLS (ordinary least squares), 278
- one vector, 22, 42
- online algorithm, 568
- online processing, 568
- Open Source Initiative, 518
- open-source, 597
- OpenMP, 616, 617
- operator matrix, 98, 263
- operator norm, 184
- optimization of vector/matrix functions, 483–499
 - constrained, 493–498
 - least squares, 487–491, 493–498
- order of a graph, 361
- order of a vector, 2
- order of a vector space, 21
- order of computations, 560
- order of convergence, 554
- order of error, 554
- order of matrix, 4
- ordinal relations among matrices, 109, 380
- ordinal relations among vectors, 23
- orthogonal array, 412
- orthogonal basis, 48–49
- orthogonal complement, 42, 144, 149
- orthogonal distance regression, 290–293, 435

- orthogonal group, 151, 343
- orthogonal matrices, binary relationship, 116
- orthogonal matrix, 149–152, 215
- orthogonal residuals, 290–293, 435
- orthogonal transformation, 215
- orthogonal vector spaces, 42, 149
- orthogonal vectors, 41
 - Exercise 2.19*., 65
- orthogonalization (Gram-Schmidt transformations), 46, 238, 582
- orthogonally diagonal factorization, 165, 181
- orthogonally diagonalizable, 165, 173, 181, 370, 375, 460
- orthogonally invariant norm, 184, 186, 188
- orthogonally similar, 165, 173, 184, 188, 259, 375
- orthonormal vectors, 41
- out-of-core algorithm, 568
- outer product, 107, 231
 - Exercise 3.16*., 204
- outer product for matrix multiplication, 587
- outer pseudoinverse, 146
- outer/inner products matrix, 389
- overdetermined linear system, 142, 241, 276
- overfitting, 288, 467
- overflow, in computer operations, 541, 545
- Overleaf, 597
- overloading, 18, 78, 185, 537, 546
- p -inverse (see also Moore-Penrose inverse), 147
- paging, 624
- parallel processing, 564, 565, 586, 588, 603, 615
- parallelogram equality, 34
- parallelotope, 87
- Parseval's identity, 49, 188
- partial ordering, 23, 109, 380
 - Exercise 8.2a*., 426
- partial pivoting, 265
- partitioned matrix, 76, 97, 149
 - determinant, 83, 140
 - sum of squares, 393, 440, 449
- partitioned matrix, inverse, 140, 149
- partitioning sum of squares, 393, 440, 449
- PBLAS (parallel BLAS), 616, 617
- PDF (probability density function), 334
- pencil, 178
- permutation, 81
- permutation matrix, 99, 105, 263, 409
- Perron root, 403, 406
- Perron theorem, 403
- Perron vector, 403, 407, 479
- Perron-Frobenius theorem, 406
- pivoting, 103, 230, 235, 265
- PLASMA, 617
- polar cone, 53
- polynomial in a matrix, 95
 - Exercise 3.26*., 205
- polynomial order, 560
- polynomial regression, 411
- polynomial, evaluation of, 570
- pooled variance-covariance matrix, 400
- population model, 479
- portability, 539, 552, 598
- Posit, 8
- positive definite matrix, 109, 119, 181–183, 240, 378–382, 460
 - summary of properties, 378–380
- positive matrix, 244, 401
- positive semidefinite matrix, 108
- positive stable, 182, 425
- power method for eigenvalues, 303–305
- precision, 531–537, 549
 - arbitrary, 524
 - double, 531, 537
 - extended, 531
 - half precision, 537
 - infinite, 524
 - multiple, 524, 549
 - single, 531, 537
- preconditioning, 271, 302, 583
 - for eigenvalue computations, 302
 - in the conjugate gradient method, 271
- primitive Markov chain, 478
- primitive matrix, 407, 478
- principal axis, 43
- principal components, 459–463
- principal components regression, 466
- principal diagonal, 70

- principal minor, 85, 122, 650
- principal submatrix, 77, 122, 227, 376, 379
 - leading, 77, 380
- probabilistic error bound, 554
- probability density function (PDF), 334
- programming model, 521, 603
- projected gradient, 494
- projected Hessian, 494
- projection (of a vector), 28, 44
- projection matrix, 388–389, 442
- projective transformation, 214
- proper value (see also eigenvalue), 153
- PSBLAS (parallel sparse BLAS), 616
- pseudo-correlation matrix, 475
- pseudoinverse (see also Moore-Penrose inverse), 147
- PV-Wave (software), 4
- Pythagorean theorem, 34
- Python, 605, 628
- Q-convergence, 566
- QL factorization, 233
- QR factorization, 232–238
 - and a generalized inverse, 236
 - computing, 617
 - matrix rank, 236
 - skinny, 233
- QR method for eigenvalues, 308–310
- quadratic convergence, 566
- quadratic form, 108, 111
- quasi-Newton method, 486
- quotient space, 130
- R (software), 8–14, 60–63, 197–202, 245–247, 293, 350–352, 499–506, 572, 605, 630–633
 - complex numbers, 13
 - computer algebra, 601
 - data frame, 501
 - merging, 502
 - tibble, 502
 - exercises, 67, 207, 249, 354, 509, 577
 - lists, 246
 - machine constants, 13
 - merging datasets, 502
 - Microsoft R Open, 633
 - missing data, 500
 - NA, 12, 500
 - Overleaf, 597
 - random numbers, 350
 - Rcpp, 632
 - RcppArmadillo, 632
 - roxygen, 631
 - RPy, 632
 - rseek, 13, 597
 - RStudio, 8, 633
 - sweave, knitr, markdown, 597
 - symbolic computation, 601
 - tibble, 502
 - views, 13, 598
- R Foundation, 518
- radix, 526
- random graph, 369
- random matrix, 337, 342–343
 - BMvN distribution, 343
 - computer generation *Exercise 7.11:*, 355
 - correlation matrix, 349
 - Haar distribution, 343
 - Exercise 7.11:*, 355
 - normal, 342
 - orthogonal *Exercise 7.11:*, 355
 - rank *Exercise 7.7:*, 353
 - Wishart, 347, 378, 458, 470
 - Exercise 7.9:*, 354
- random number generation, 11–12, 348–352
 - random matrices *Exercise 7.11:*, 355
- random variable, 334
- range of a matrix, 69
- rank deficiency, 119, 162
- rank determination, 590
- rank of a matrix, 117–139, 236, 469, 590
 - of idempotent matrix, 383
 - rank-revealing QR, 236, 469
 - statistical tests, 469–473
- rank of an array, 3
- rank reduction, 590
- rank($\cdot$), 117
- rank, linear independence, 117, 590
- rank, number of dimensions, 3
- rank-one decomposition, 181
- rank-one update, 220, 274
- rank-revealing QR, 236, 469, 590
- rate constant, 566
- rate of convergence, 566
- rational fraction, 549

- Rayleigh quotient, 97, 175, 424, 496
- Rcpp, 632
- RcppBlaze, 617
- RCR (Replicated Computational Results), 611
- real numbers, 525
- real-time algorithm, 568
- recursion, 569
- reduced gradient, 494
- reduced Hessian, 494
- reduced rank regression problem, 470
- reducibility, 303, 366–368, 404
 - Markov chains, 478
- reflection, 217–219
- reflector, 219
- reflexive generalized inverse, 146
- register, in computer processor, 542
- regression, 430–459, 463–468
 - nonlinear, 487
 - variable selection, 464
- regular graph, 361
- regular matrix (see also diagonalizable matrix), 167
- regularization, 287, 434, 466
- relative error, 541, 552, 584
- relative spacing, 528
- Reliable Computing*, 550
- Replicated Computational Results (RCR), 611
- Reproducible R Toolkit, 633
- reproducible research, 610–611, 633
- residue arithmetic, 550
- restarting, 583
- reverse communication, 623
- $\rho(\cdot)$ (spectral radius), 160
- Richardson extrapolation, 568
- ridge regression, 260, 393, 435, 447, 466
 - Exercise 9.7a*, 508
- right direct product, 112
- right inverse, 126
- robustness (algorithm or software), 557
- root of a function, 544
- root-free Cholesky, 241
- Rosser test matrix, 609
- rotation, 215–218, 222
- rounding, 531
 - rounding error, 544, 552
- row echelon form, 131
- row rank, 117
- row space, 70
- row-major, 580, 598, 604
- row-sum norm, 185
- roxygen, 631
- RPy, 632
- RQ factorization, 233
- RStudio, 8, 633
- Samelson inverse, 39
- sample variance, computing, 558
- saxpy**, 18
- scalability, 563
- ScaLAPACK, 616, 617
- scalar, 17
- scalar product, 31
- scaled matrix, 397
- scaled vector, 58
- scaling of a vector or matrix, 259
- scaling of an algorithm, 560, 563
- Schatten p norm, 189
- Schur complement, 139, 440, 458
- Schur factorization, 166–167
- Schur norm (see also Frobenius norm), 187
- SDP (semidefinite programming), 378
- Seidel adjacency matrix, 364
- self-adjoint matrix (see also Hermitian matrix), 70
- semidefinite programming (SDP), 378
- seminorm, 33
- semisimple eigenvalue, 163, 168
- sequences of matrices, 191
- sequences of vectors, 40
- shape of matrix, 4
- shearing transformation, 214
- Sherman-Morrison formula, 275, 445
- shifting eigenvalues, 302
- shrinkage, 435
- side effect, 612
- $\sigma(\cdot)$ (sign of permutation), 81, 105
- $\sigma(\cdot)$ (spectrum of matrix), 159
- sign bit, 523
- sign($\cdot$), 22
- significand, 526
- similar canonical form, 167
- similar matrices, 164
- similarity matrix, 401
- similarity transformation, 164–167, 305, 309

- simple eigenvalue, 163
- simple graph, 361
- simple matrix (see also diagonalizable matrix), 167
- single precision, 531, 537
- singular matrix, 119
- singular multivariate normal distribution, 340, 470
- singular value, 179, 463, 590
 - relation to eigenvalue, 180
- singular value decomposition, 178–181, 312, 368, 462, 590
 - computing, 312
 - uniqueness, 180
- size of matrix, 4
- skew diagonal element, 74
- skew diagonal matrix, 71
- skew symmetric matrix, 70, 74
- skew upper triangular matrix, 72, 420
- skinny QR factorization, 233
- smoothing matrix, 393, 447
- software testing, 607–609
- SOR (method), 268
- span($\cdot$), 21, 29, 69
- spanning set, 21, 29
 - of a cone, 52
- Spark, 518, 522
- Spark (software system), 603
- sparse matrix, 73, 245, 267, 580, 584, 615, 616
 - index-index-value, 607
 - software, 616
 - CUDA, 619
 - storage mode, 607
- spectral circle, 160
- spectral condition number, 258, 259, 279
- spectral decomposition, 173, 181
- spectral norm, 186, 189
- spectral projector, 174
- spectral radius, 160, 186, 190, 268
- spectrum of a graph, 424
- spectrum of a matrix, 159–164
- splitting extrapolation, 568
- square root matrix, 183, 239, 240, 377
- stability, 266, 557
- standard deviation, 58, 396
 - computing the standard deviation, 558
- Standard Template Library, 628
- standards (see also specific standard), 518
- stationary point of vector/matrix functions, 485
- statistical reference datasets (StRD), 609
- steepest descent, 484, 486
- Stiefel manifold, 151
- stiff data, 559
- stochastic matrix, 409
- stochastic process, 476–499
- stopping criterion, 566
- storage mode, for matrices, 605–607
- storage unit, 522, 526, 538
- Strassen algorithm, 587–588
- StRD (statistical reference datasets), 609
- stride, 580, 598, 613
- string, character, 520
- strongly connected graph, 366
- submatrix, 76, 97
- subspace of vector space, 24
- successive overrelaxation, 268
- summation, 542, 543
- summing vector, 42
- Sun ONE Studio Fortran 95, 550
- superlinear convergence, 566
- SVD (singular value decomposition), 178–181, 312, 368, 462, 590
 - computing, 312
 - uniqueness, 180
- sweep operator, 440
- Sylvester equation, 114
- Sylvester’s law of nullity, 135
- symbolic computations, 601
- symmetric matrix, 70, 74, 132, 172–174, 182–183, 370–375
 - eigenvalues/vectors, 172–174, 182–183
 - equivalent forms, 132
 - inverse of, 138
 - summary of properties, 370
- symmetric pair, 311
- symmetric storage mode, 76, 606
- Taylor series, 322, 485
- Template Numerical Toolkit, 627
- tensor, 3

- term-document matrix, 368
- test problems for algorithms or software, 584, 607–609
 - Exercise 3.46*., 208
 - consistency test, 584, 609
 - Ericksen matrix, 608
 - Exercise 12.8*., 635
 - Hilbert matrix, 607
 - Matrix Market, 609
 - randomly generated data, 607
 - Exercise 11.5*., 594
 - Rosser matrix, 609
 - StRD (statistical reference datasets), 609
 - Wilkinson matrix, 608
 - Exercise 12.9*., 635
 - Wilkinson's polynomial, 556
- testable hypothesis, 456
- testing software, 607–609
- thread, 603
- Tikhonov regularization, 288, 466
- time series, 481–499
 - variance-covariance matrix, 415, 482
- Toeplitz matrix, 414, 482
 - circulant matrix, 415
 - inverse of, 415
 - Exercise 8.12*., 427
 - Exercise 12.12*., 636
- total least squares, 290, 435
- $\text{tr}(\cdot)$, 80
- trace, 80
 - derivative of, 330
 - of Cayley product, 106
 - of idempotent matrix, 383
 - of inner product, 109
 - of Kronecker product, 114
 - of matrix inner product, 116
 - of outer product, 109
 - relation to eigenvalues, 159
- trace norm, 189
- transition matrix, 477
- translation transformation, 218
- transpose, 73
 - determinant of, 82
 - generalized inverse of, 142
 - inverse of, 125
 - norm of, 184
 - of a vector, 96
 - of Cayley product of matrices, 92
 - of Kronecker product, 113
 - of partitioned matrices, 78
 - of sum of matrices, 78
 - trace of, 80
- trapezoidal matrix, 71, 227, 233
- triangle inequality, 33, 184
 - Exercise 2.12b*., 64
- triangular matrix, 71, 102, 227
 - determinant of, 83
 - inverse of, 139
 - multiplication, 95
- tridiagonal matrix, 72
- triple scalar product *Exercise 2.21c*., 66
- triple vector product *Exercise 2.21d*., 66
- truncation error, 50, 117, 555, 556
- twos-complement representation, 523, 540
- type 2 matrix, 72, 415
- ulp (“unit in the last place”), 530
- underdetermined linear system, 141
- underflow, in computer operations, 528, 545
- Unicode, 519
- union of vector spaces, 25
- unipotent matrix, 95
- unit in the last place (ulp), 530
- unit roundoff, 529
- unit triangular matrix, 95
- unit vector, 22, 29, 43, 93
- unitarily diagonalizable, 165, 375, 419
- unitarily similar, 165
- unitary matrix, 150
- unrolling do-loop, 624
- updating a solution, 274, 283, 444–447
 - regression computations, 444–447
- upper Hessenberg form, 73, 308
- upper triangular matrix, 71
- usual norm (see also Frobenius norm), 187
- $V(\cdot)$ (variance operator), 58, 336
- $\mathcal{V}(\cdot)$ (vector space), 29, 69
- Vandermonde matrix, 411
 - Fourier matrix, 416
- variable metric method, 486
- variable selection, 464
- variance, computing, 558
- variance-covariance matrix, 336, 343

- Kronecker structure, 114, 343, 458
- positive definite approximation, 474
- sample, 396, 460
- $\text{vec}(\cdot)$, 75
- vec-permutation matrix, 100
- $\text{vecdiag}(\cdot)$, 74
- $\text{vech}(\cdot)$, 76
- vector, 2
 - centered vector, 57
 - mean vector, 42
 - normal vector, 42
 - normalized vector, 38
 - null vector, 22
 - one vector, 22, 42
 - “row vector”, 96
 - scaled vector, 58
 - sign vector, 22
 - summing vector, 22, 42
 - transpose, 96
 - unit vector, 22
 - zero vector, 22
- vector derivative, 317–330
- vector processing, 615
- vector space, 20–30, 69, 79, 144, 145
 - basis, 29–30
 - definition, 20
 - dimension, 21
 - direct product, 28
 - direct sum, 25–27
 - direct sum decomposition, 27
 - essentially disjoint, 24
 - generated by a matrix, 69
 - generated by a set of vectors, 21
 - intersection, 25
 - null vector space, 20
 - of matrices, 79
 - order, 21
 - set operations, 23
 - subspace, 24
 - union, 25
- vector subspace, 24
- vectorized processor, 565
- vertex of a graph, 361
- volume as a determinant, 86, 340
- weighted graph, 361
- weighted least squares, 443
 - with equality constraints *Exercise 9.3d*, 507
- weighted norm, 36, 111
- Wilk’s Λ , 458
- Wilkinson matrix, 608
- Wishart distribution, 378
 - Exercise 7.9*, 354
- Woodbury formula, 275, 445
- word, computer, 522, 526, 538
- XDR (external data representation), 539
- Yule-Walker equations, 483
- Z-matrix, 425
- zero matrix, 93, 117
- zero of a function, 544
- zero vector, 22